

# What Is Attacking My Pine? The Case of the Deodar Weevil

Many pine owners are able to identify the damage caused by southern pine beetles (*Dendroctonus frontalis*) (SPB) or Ips engraver beetles (*Ips* spp.). Many, however, have never heard of the deodar weevil (*Pissodes nemorensis*) unless they have experienced an outbreak on their property. Fortunately, this native insect leaves very distinctive signs and tree symptoms that landowners can use to survey their pines for this pest.

Deodar weevils are native insects that attack a variety of conifers, such as all loblolly (*Pinus taeda*), longleaf (*P. palustris*), shortleaf (*P. echinata*), and slash (*P. elliottii*). In response to deodar weevils and other insects, pines have developed natural defenses which effectively prevent attacks when trees are healthy and growing rapidly. During most years, deodar weevils only attack a few suppressed, unhealthy trees scattered in well-managed plantations. As such, the deodar weevil is not an insect that regularly alters traditional pine management.

Pine owners should be mindful of the risk deodar weevils can pose under certain environmental conditions. Being able to differentiate deodar damage from that of other insect pests is important for proper stand management. Rapid identification of a deodar outbreak can help guide post-attack management treatments and reduce future damage. The goal of this publication is to help forest landowners identify deodar weevils and develop a plan to prevent or moderate damage to pine plantations.

## Conditions Favoring Outbreaks

Tree stress is an important factor in deodar weevil outbreaks. Deodar outbreaks during late summer droughts can be as devastating as those caused by Ips engraver beetles; however, droughts rarely lead to outbreaks because deodar weevils also require favorable breeding conditions (warm fall temperatures) the previous year. Nevertheless, when this combination of factors occurs, deodars, along with Ips engraver beetles, are capable of killing large numbers of stressed pines (Figure 1). Factors besides drought that increase the risk of outbreak by inducing tree stress include overly stocked pine stands and wind, hail, or logging damage the preceding fall. Stands heavily fertilized with nitrogen or located near chicken houses also seem to attract deodars.

## Associations with Pitch Canker

Deodar weevils are vectors of pitch canker (*Fusarium moniliforme*). Although deodars are capable of disseminating



**Figure 1. An outbreak of deodar weevils and Ips engraver beetles occurred in 2015. Together, these pests contributed to widespread pine mortality in central Mississippi (brown tops). Landsat image via Google Maps, Copeiah County, 2016.**

pitch canker in all seasons, the majority of spread occurs during the fall when fungus-carrying weevils bore into trees to reproduce. It is important to note that not all deodar weevils carry the fungus.

Deodars must first pick up the fungus from an infected tree before it can spread to a healthy pine. Pitch canker kills a small area of stem or branch, causing it to exude large amounts of sap or pitch. As such, pitch running down stems and branches is a key symptom of pitch canker (Figure 2). The combination of pitch canker and deodar weevils can be lethal in pine plantations with reports of 40 percent top dieback and 30 percent mortality occurring in a single year.

## Life Cycle

The life cycle of deodar weevils begins in the spring (April to May) with adults inside wood chip cocoons under pine stem bark (Figure 3). Adults (Figure 4) bore out of the tree and begin feeding on succulent tops and lateral branches of pine trees (Figure 5). Adult activity is diminished during summer heat. Adults become active again in cooler fall temperatures. Fall is the peak time for adult feeding and breeding. Deodars prefer stressed, dying, or dead trees for breeding. Often, they



**Figure 2.** Sap running down a pine stem or branch is a symptom of pitch canker. Under the bark are pockets of pitch. One potential vector of this disease is the deodar weevil. Photos by Stephen Dicke, Leake County, 2012.


seek trees previously attacked by SPB or Ips. Slash can also be used for reproduction. Female adults puncture the bark with their snouts and typically lay between one and four eggs inside each feeding puncture. Legless, white, grub-like larvae with brown heads hatch and mine random paths under bark over the course of fall and winter. It is this larval feeding that ultimately can girdle and kill the tree (Figure 6). Trees dying in the winter may indicate a heavy deodar infestation. By February to March, larvae have grown to one-fourth of an inch long. Larvae then surround themselves with half-inch, football-shaped wood chip cocoons. By April, larvae pupating inside chip cocoons slowly change into adults (Figure 7). This month-long pupation period is when deodar weevils are most vulnerable to predation. The pupation period also gives landowners an opportunity to identify and control this pest.

## Deodar Weevil Identification

Like all weevils, adult deodar weevils are easy to distinguish from other wood-boring insects. Look for their long snouts with antennae (Figure 4). Body color ranges from gray to reddish-brown, with white spots for camouflage. Deodar weevil adults average a quarter-inch length, much larger than Ips engraver beetles or SPB and similar in size to the black turpentine beetle (*Dendroctonus terebrans*).



**Figure 3.** (top) A round hole in a chip cocoon indicates that an adult deodar weevil has successfully emerged. (bottom) Bark sloughed off a heavily infested stem exposes a high number of wood chip cocoons. Photos by Stephen Dicke, Waltham and Rankin Counties, 2016.



**Figure 4.** An adult deodar weevil. Like all weevils, deodars have long snouts with antennae. Weevils can vary from red to black with splatches of camouflage. Photo by Natasha Wright, Cook's Pest Control, Bugwood.org.

## Signs and Symptoms of Deodar Weevils

### Fading Tree Crowns and Dying Tops

Fading crowns in the fall or winter are a common symptom of deodar weevil activity (Figure 5); however, this symptom alone does not confirm presence of a deodar infestation—or any other insect, for that matter. Instead, crown fade simply indicates stress, which can be caused by a variety of factors. Following fading, pine tops can die, and, ultimately, the stem can die if damage is great enough (Figure 1). Crowns fading in late spring and summer are most likely caused by something other than deodar weevils.

### Chip Cocoon

The most definitive sign of a deodar infestation is the chip cocoon (Figure 3). Created by larvae in preparation of their pupation, cocoons are visible as early as February. Deodar weevils tend to occupy the bottom 10 feet of the main stem, so cocoons are relatively easy to spot. Bark can also slip off heavily infested stems, exposing the chip cocoons. Intact



**Figure 5. (left)** Adult deodar weevil feeding activities on pine tops and branches can initially cause foliage decline and yellowing. This can lead to tip dieback and, ultimately, tree death. Photo by Stephen Dicke, Adams County, 2015. **(right)** A symptom of adult deodar weevil feeding damage in pine tops is a general lack of foliage in the fall to winter. Photo by Stephen Dicke, Leake County, 2012.

cocoons mean larvae and pupae are still inside. There is still time to control this population of weevils.

Once cocoons are found, pay close attention to the existence of an exit hole and its shape. Cocoons with perfectly round exit holes indicate a successful adult deodar emergence (Figure 3). Imperfect exit holes are a good sign for landowners. Cocoons that appear to be ripped open are indicative of woodpeckers (*Picidae* spp.) preying on larvae and pupae (Figure 8). Birds are an important natural control agent of deodar populations.

## Deodar Weevil Risk

Loblolly pine is the most common host for deodar weevils, but all pine species in Mississippi can be attacked. Pure stands of pine are at greater risk of deodar weevil outbreaks than mixed stands of pine-hardwood. The risk of attack by deodar weevils begins about age 5, when stems are large enough to support larval galleries. Pines at that age are also very sensitive to mortality from crown damage. Risk of attack peaks for pine plantations before the first thin, when resource competition begins to reduce tree vigor. After the first thinning, the risk of a deodar attack drops. Carefully scout your pine plantation following summer to fall droughts, wind and hail storms, and fall logging damage. Also, scout pine plantations adjacent to chicken houses and in plantations heavily fertilized with nitrogen.

Finding some signs of deodar weevils is not of great concern. This is because deodars do not kill healthy pine trees. Thus, immediate action is rarely required.

## Preemptive Management

The best way to prevent a deodar outbreak is to maintain a vigorously growing plantation. Rapidly growing pines are typically healthy and will have ample resin flow to cover and



**Figure 6. Galleries created by deodar weevil larvae in the winter. Tunnels are much larger than those of *Ips* engraver and southern pine beetles. Photo by Stephen Dicke, Leake County, 2016.**

kill newly hatched larvae. For this reason, adult beetles do not usually lay their eggs on healthy pines.

It is perfectly normal to have suppressed, unhealthy trees scattered throughout a plantation before a thinning. Landowners should not be concerned if deodar signs are found on these suppressed trees. In fact, suppressed tree mortality may actually be beneficial, as it releases better trees from competition. In addition, unlike SPB or *Ips* engraver beetles, deodar infestations will not actively spread throughout the growing season.



**Figure 7. Deodar weevil larvae form a wood chip cocoon in late winter. The middle cocoon was opened to reveal a pupa. Termites left the wood chip cocoons alone. Photo by Stephen Dicke, Copiah County, 2016.**

Landowners should focus on thinning to maintain tree vigor. As a general rule of thumb, landowners should schedule thinning once tree diameter growth falls below an annual threshold of 5 percent. Egg-laying females usually avoid pines growing at 5 percent or greater. Thus, thinning will help keep your stand safe in the event of a deodar outbreak.

One factor landowners should be aware of is residual damage after thinning. Trees damaged in thinning operations typically become stressed, increasing the risk of a deodar attack. Fortunately, much of this risk can be overcome by thinning either before or after the peak breeding season (September to November). For more thinning guidance, refer to MSU Extension [Publication 2260 Are My Pines Ready to Thin?](#) or [Publication 3664 Simplifying Southern Pine Thinning with Stand Density Index](#), and download the MSUES Pine Thin app.

A sound strategy for preventing deodar damage is to plant pines on sites where they are best suited to grow. Compared to most hardwood species, pines have a relatively wide range of sites on which they can be planted. However, even pines have their limits. Planting a pine on the wrong site increases stress and decreases vigor. For more information on pine site considerations, refer to Extension [Publication 1776 Planting Southern Pines: A Guide to Species Selection and Planting Techniques](#).



**Figure 8. The shattering of cocoons is a sign of woodpecker (*Picidae* spp.) predation on deodar weevil pupae. Photo by Stephen Dicke, Rankin County, 2016.**

## Summary

Although deodar weevil outbreaks are rare, it is important to remember the destructive potential of this insect. This is particularly true for young pine plantations, where risks are elevated. In most years, deodar weevils pose only a minor threat to pine plantations. This is because healthy pines are able to resist attack. Also, deodar weevils have only one generation per year, reducing potential for an expanding attack; therefore, signs of deodar attacks on suppressed pines is not a major concern. Stresses that can lead to deodar outbreaks include climatic events such as droughts, wind and hail storms, logging damage, and ammonia gas from chicken houses. Scouting young pine plantations following these stressors can help identify deodar attacks. Outbreaks are still hard to predict. As such, the best way to defend against an outbreak is to maintain stand vigor. This includes planting trees on appropriate growing sites and thinning before the onset of intense competition. These basic steps will help keep pines safe.

**Publication 3057** (POD-05-26)

Reviewed by **Kristy McAndrew**, PhD, Assistant Professor, Forestry, from an earlier revision by A. Brady Self, Extension Professor, Forestry. Written by John Willis, PhD, former Assistant Professor; Stephen Dicke, PhD, Professor Emeritus, Forestry; and John Riggins, PhD, former Professor, Entomology.

Copyright 2026 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.

Produced by Agricultural Communications.

Mississippi State University is an equal opportunity institution. Discrimination is prohibited in university employment, programs, or activities based on race, color, ethnicity, sex, pregnancy, religion, national origin, disability, age, sexual orientation, genetic information, status as a U.S. veteran, or any other status to the extent protected by applicable law. Questions about equal opportunity programs or compliance should be directed to the [Office of Civil Rights Compliance](#), 231 Famous Maroon Band Street, P.O. 6044, Mississippi State, MS 39762.

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. ANGUS L. CATCHOT JR., Director